

April 2020 ~ Resource #360423

COVID-19 and Pharmacotherapy

The **first chart below** provides information or resources on pharmacotherapy of interest for COVID-19, the disease caused by the SARS-CoV-2 virus. Additional resources on pharmacotherapy, which are frequently updated, include:

- The **American Society of Health-System Pharmacists** evidence table of COVID-19 treatments (<https://www.ashp.org/-/media/assets/pharmacy-practice/resource-centers/Coronavirus/docs/ASHP-COVID-19-Evidence-Table>).
- The **British Columbia Ministry of Health** evidence review (http://www.bccdc.ca/Health-Professionals-Site/Documents/Guidelines_Unproven_Therapies_COVID-19.pdf).

At this point, no pharmacotherapy has been proven effective for COVID-19, so treatment is largely supportive. Resources pertinent to supportive therapy include:

- The **WHO** guidance for the treatment of suspected COVID-19 severe acute respiratory infection ([https://www.who.int/publications-detail/clinical-management-of-severe-acute-respiratory-infection-when-novel-coronavirus-\(ncov\)-infection-is-suspected](https://www.who.int/publications-detail/clinical-management-of-severe-acute-respiratory-infection-when-novel-coronavirus-(ncov)-infection-is-suspected)).
- The **Surviving Sepsis Campaign** COVID-19 guidelines (<https://sccm.org/SurvivingSepsisCampaign/Guidelines/COVID-19>).
- The **NIH** general treatment guidelines (<https://covid19treatmentguidelines.nih.gov/>).

The **second chart below** addresses common questions about pharmacotherapy as it relates to COVID-19.

****Search www.clinicaltrials.gov for the latest information on COVID-19 clinical trials.****

Abbreviations: ACE = angiotensin-converting enzyme; ARB = angiotensin receptor blocker; ARDS = acute respiratory distress syndrome; COPD = chronic obstructive pulmonary disease; CT = computed tomography; ECMO = extracorporeal membrane oxygenation; EUA = Emergency Use Authorization; IDSA = Infectious Diseases Society of America; IL = interleukin; NIH = National Institutes of Health; NSAIDs = nonsteroidal anti-inflammatory drugs; SARS = severe acute respiratory syndrome; SARS-CoV-2 = the virus that causes COVID-19 disease; tPA = tissue plasminogen activator; TNF = tumor necrosis factor; WHO = World Health Organization

TREATMENTS OF INTEREST

Drug	Pertinent Information or Resources
Anakinra (<i>Kineret</i>)	<p>Note that DOSES provided are examples only for ADULTS; the optimal dose has not been determined for any treatment.</p> <ul style="list-style-type: none">• Anakinra is an IL-1 antagonist. IL-1 may have a role in ARDS.⁶⁵• Anakinra 5 mg/kg twice daily intravenously in moderate to severe ARDS (non-ventilator) and inflammation (elevated C-reactive protein and/or ferritin) (n=29) was associated with improved survival compared to a similar historical cohort (90% vs 56%, p = 0.009).⁶⁵ These patients also received hydroxychloroquine and lopinavir/ritonavir.⁶⁵ A lower dose of anakinra (100 mg twice daily subcutaneously) did not seem to provide benefit.⁶⁵

More . . .

Drug	Pertinent Information or Resources
Azithromycin	<p>Note that DOSES provided are examples only for ADULTS; the optimal dose has not been determined for any treatment.</p> <ul style="list-style-type: none"> • Macrolides have <i>in vitro</i> antiviral (e.g., Zika, Ebola), anti-inflammatory, and immunomodulatory activity.^{2,7} • Insufficient evidence to support widespread use [Evidence level C].^{2,28} • Was used in a small, widely publicized study with hydroxychloroquine in six patients to prevent bacterial superinfection in COVID-19 patients (See hydroxychloroquine, below).² Subsequent observational data including 74 additional patients suggests that the combination can reduce viral load and perhaps improve the clinical course, but there was no comparator group.²⁸ Also see the hydroxychloroquine section below for information on its use in a U.S. cohort study.⁷⁵ • NIH guidelines recommend against the use of azithromycin plus hydroxychloroquine outside of a clinical trial.⁵⁰ • Studies for COVID-19 treatment include various dosing regimens (usually azithromycin 500 mg x 1 then 250 mg once daily for four days) WITH chloroquine, hydroxychloroquine, or other antimicrobials. See www.clinicaltrials.gov for the latest information on these studies. • When used with hydroxychloroquine or chloroquine (and other QT prolonging medications), QT prolongation is of increased concern.^{2,6}
Aviptadil	<ul style="list-style-type: none"> • Investigational synthetic form of vasoactive intestinal polypeptide. Has anti-IL-6 and anti-TNF activity. Phase I trial suggests benefit in ARDS. No COVID-19 data. • Clinical trial is planned for COVID-19-associated ARDS. See www.clinicaltrials.gov for more information.
Baloxavir (Xofluza)	<ul style="list-style-type: none"> • No COVID-19 data.
Chloroquine phosphate* *Chloroquine phosphate 500 mg = chloroquine base 300 mg ⁶ <i>Continued...</i>	<p>Efficacy</p> <ul style="list-style-type: none"> • Inhibits SARS-CoV-2 <i>in vitro</i>,² but clinical trials have not shown benefit against other viruses.¹⁸ Also has immunomodulating effects.²⁶ Early reports suggested that for COVID-19 pneumonia, it could speed clinical improvement and viral clearance.³ • The FDA has revoked its EUA for chloroquine because it is unlikely to be effective, based on data from the EUA and elsewhere.⁷³ • Clinical trials are planned on the use of chloroquine to prevent COVID-19 in healthcare workers. • See www.clinicaltrials.gov for regimens being studied. <p>Dosing</p> <ul style="list-style-type: none"> • The FDA suggested, as part of their now-revoked EUA for chloroquine, for patients weighing ≥ 50 kg, a chloroquine phosphate dose of 1 g on day one, followed by 500 mg once daily for four to seven days.⁴ This suggested dosing regimen is unlikely to produce an antiviral effect.⁷³

More...

Drug	Pertinent Information or Resources Note that DOSES provided are examples only for ADULTS ; the optimal dose has not been determined for any treatment.
Chloroquine, continued	<p>Safety</p> <ul style="list-style-type: none"> In addition to efficacy concerns, the FDA’s revocation of its EUA for chloroquine was based on adverse effects; its known and potential benefits no longer outweigh the known and potential side effects (e.g., serious cardiac events and other serious side effects).³³ The FDA recommends against chloroquine use for COVID-19 outside of a clinical trial.³³ Adverse effects are not well-characterized at the doses studied for COVID-19. In general, potential adverse effects include: gastrointestinal side effects (take with food or milk), headache, hypoglycemia, QT prolongation and other conduction disturbances (especially with hypokalemia, hypomagnesemia, or heart disease), cardiomyopathy, myopathy, movement disorders, neurotoxicity, ocular toxicity, ototoxicity, anemia, thrombocytopenia, neutropenia, bone marrow suppression, serious dermatologic reactions, and psoriasis flare.^{4,27} Monitor electrolytes, glucose, complete blood count, electrocardiogram, baseline renal and hepatic function, knee and ankle reflexes, vision, and mental status.^{4,6,27} A Brazilian study of chloroquine phosphate 600 mg twice daily vs 450 mg twice daily stopped the high-dose arm due to higher instance of QT prolongation >500 milliseconds (18.9% vs 11.1%) and mortality (39% vs 15%).⁴¹ All patients received azithromycin.⁴¹ When used with azithromycin (and other QT-prolonging medications), QT prolongation is of increased concern.^{2,4,6}
Colchicine	<ul style="list-style-type: none"> Based on its anti-inflammatory effect, there is interest in using colchicine to alter the clinical course of COVID-19 in both inpatients and higher-risk outpatients. The open-label GRECCO-19 study randomized patients to colchicine plus standard care or standard care (n = 105). The clinical primary endpoint, which included measurements of inflammation and clinical deterioration, occurred in 14% of the control group vs 1.8% in the colchicine group (p=0.02).⁹ This study’s findings are considered “hypothesis-generating” only.⁹ Additional clinical trials are underway. See www.clinicaltrials.gov for more information. Keep in mind colchicine’s toxicities and drug interactions. See our chart, <i>Colchicine Dosing and Interactions</i>, for details.
Convalescent Plasma (COVID-19) <i>Continued...</i>	<ul style="list-style-type: none"> No large studies have been published, but small case series in patients hospitalized with severe COVID-19 show promise (e.g., defervescence, radiographic improvement, improved oxygen support requirements, viral clearance, improved clinical condition).⁶²⁻⁶⁴ It appears well-tolerated.⁶²⁻⁶⁴ Concerns include allergic reaction, viral infection, and increased clotting risk.^{70,71} There are three pathways for administering or studying COVID-19 convalescent plasma: clinical trials (www.clinicaltrials.gov or https://covidcp.org/), expanded access (https://uscovidplasma.org/), or single patient emergency IND. The FDA has guidance for use of convalescent plasma at https://www.fda.gov/vaccines-blood-biologics/investigational-new-drug-ind-or-device-exemption-ide-process-cber/recommendations-investigational-covid-19-convalescent-plasma.

More...

Drug	Pertinent Information or Resources
Convalescent plasma, continued	<p>Note that DOSES provided are examples only for ADULTS; the optimal dose has not been determined for any treatment.</p> <ul style="list-style-type: none"> See https://www.ccpp19.org/ or https://uscovidplasma.org to find out how recovered patients can donate their plasma.
Corticosteroids	<ul style="list-style-type: none"> There is concern that corticosteroids have the potential to delay viral clearance, as observed in MERS-CoV and SARS.¹⁰ However, they are being studied for COVID-19. In one institution in China, methylprednisolone use in patients with COVID-19 ARDS was associated with reduced mortality.¹⁶ This and other cohort studies were limited by confounding, and inclusion of patients with various disease severities and concomitant treatments.⁴⁶ Data from the open-label RECOVERY trial, in which 2,104 patients were randomized to oral or intravenous dexamethasone 6 mg/day for 10 days, suggests a mortality benefit for COVID-19 patients requiring oxygen, especially for those requiring ventilation, over usual care (n = 4,321).³¹ NNT = 8 to prevent one death in ventilated patients, or 34 in patients requiring oxygen but not ventilation. It did not provide a mortality benefit (and may harm) patients not requiring oxygen. It also did not provide a mortality benefit for early disease (symptoms for a week or less). This suggests that dexamethasone's mechanism involves an anti-inflammatory effect rather than an antiviral effect, because inflammation is more common in advanced disease, while viral replication is at maximum in early disease. The IDSA suggests dexamethasone 6 mg/day x 10 days (or until discharge, if earlier), for patients hospitalized with severe COVID-19 (oxygen saturation $\leq 94\%$ on room air; need for supplementation oxygen, mechanical ventilation, or extracorporeal membrane oxygenation). If dexamethasone is not available, methylprednisolone 32 mg or prednisone 40 mg daily can be used.⁴⁶ NIH guidelines similarly recommend dexamethasone 6 mg/day x 10 days in COVID-19 patients who require oxygen or mechanical ventilation.⁵⁰ Corticosteroids are not recommended for COVID-19 patients not requiring treatment with supplemental oxygen.^{46,50} Harms of corticosteroids include hyperglycemia, agitation, confusion, and infection risk.⁴⁶ WHO Guidelines recommend that at this time, outside of clinical trials, corticosteroids should be reserved for patients with specific indications for them (e.g., sepsis, COPD, asthma, septic shock, ARDS) with consideration to risk vs benefit.¹⁰ These guidelines predate the RECOVERY trial. Ciclesonide (<i>Alvesco</i>) and inhaled budesonide are being studied for treatment of COVID-19. See www.clinicaltrials.gov for more information.
Dapagliflozin	<ul style="list-style-type: none"> No data. Dapagliflozin is being studied in COVID-19 patients with respiratory failure and with hypertension, diabetes, heart disease, or advanced renal disease to prevent organ failure, based on its known renal and cardiac benefit (DARE-19 study). See www.clinicaltrials.gov for more information.

More . . .

Drug	Pertinent Information or Resources
Famotidine	<p>Note that DOSES provided are examples only for ADULTS; the optimal dose has not been determined for any treatment.</p> <ul style="list-style-type: none"> • Interest in famotidine as a COVID-19 treatment stems from observations in China that patients who were taking famotidine who were infected with COVID-19 had better outcomes.⁵⁵ • In a retrospective U.S. study (n = 1,620), famotidine use (10 to 40 mg/day; n = 84) within 24 hours of admission was associated with reduced risk of death or intubation in hospitalized COVID-19 patients.⁶⁷ • The IDSA suggests against use of famotidine for COVID-19 outside of a clinical trial.⁴⁶ See www.clinicaltrials.gov for more information.
Hydroxy-chloroquine	<p>Efficacy</p> <ul style="list-style-type: none"> • Is a more potent inhibitor of SARS-CoV-2 than chloroquine <i>in vitro</i>.² Also has immunomodulating effects.²⁷ • The FDA has revoked its EUA for hydroxychloroquine because it is unlikely to be effective, based on data from the EUA and elsewhere.⁷³ • The hydroxychloroquine arm of the large RECORD study was also stopped due to lack of efficacy.³¹ • Early enthusiasm for hydroxychloroquine was based on a widely publicized open-label, randomized study in hospitalized patients testing positive for SARS-CoV-2.² Six of 26 hydroxychloroquine patients were lost to follow-up: one due to death, three due to intensive care admission, one due to side effects (nausea), and one who left the hospital. Viral clearance at day six was 70% in the 20 remaining hydroxychloroquine patients vs 12.5% of the control patients (n = 16).² Six treated patients also received azithromycin 500 mg on day one, then 250 mg on days two through five to prevent bacterial infection.² In the combination group, viral clearance was 100% at day six vs 57.1% in the hydroxychloroquine-alone group.² Also see subsequent observational data under “Azithromycin,” above. • In a pilot study in China, 30 patients were randomized to hydroxychloroquine 400 mg/day (it is unclear if this was divided) for five days, or usual care. There was no difference between groups in viral clearance at day seven, length of stay, or time to defervescence.²⁹ In a study of 62 hospitalized patients with mild disease, 31 patients were randomized to hydroxychloroquine 200 mg twice daily. Time to recovery (defervescence and cough remission) was shortened by about one day in the treatment group. On day six, pneumonia was improved per CT in more patients in the treatment group. Four patients progressed to severe disease, all in the control group.³⁹ A subsequent open-label Chinese study (n=150) randomized patients to usual care or hydroxychloroquine 1,200 mg/day for three days, then 800 mg/day for two (mild to moderate disease) or three weeks (severe disease).⁴² Viral clearance was similar on days 4, 7, 10, 14, 21, and 28 for the two groups. Hydroxychloroquine did not seem to hasten symptom improvement. Thirty percent of hydroxychloroquine patients had adverse effects. • A retrospective French cohort study found no benefit of early hydroxychloroquine administration to 84 hospitalized patients in regard to need for intensive care, or mortality.⁴³ • Similarly, in a retrospective U.S. Veterans Affairs study (n=368), hydroxychloroquine, alone or with azithromycin, was not associated with reduced need for mechanical ventilation in hospitalized patients. Mortality was higher in patients <p><i>Continued...</i></p>

More. . .

Drug	Pertinent Information or Resources
Hydroxy-chloroquine, continued	<p>Note that DOSES provided are examples only for ADULTS; the optimal dose has not been determined for any treatment.</p> <p>who received hydroxychloroquine alone vs no hydroxychloroquine.⁴⁹</p> <ul style="list-style-type: none"> • Other retrospective U.S. studies suggest no benefit. In one (n = 1,376), patients received hydroxychloroquine 600 mg twice daily on day 1, then 400 mg daily for a median of five days. Some patients also received azithromycin or sarilumab. About half of patients began treatment within 24 hours of presentation. Hydroxychloroquine use was not associated with a reduced risk of a composite outcome of death or intubation.⁶⁰ In another study (n=1,438), neither hydroxychloroquine alone, azithromycin alone, nor the combination was associated with improved in-hospital mortality, but the combination was associated with cardiac arrest.⁶⁶ • One large (n = 2,541) retrospective U.S. cohort study found reduced mortality with hydroxychloroquine +/- azithromycin vs usual care.⁷⁵ Patients received 400 mg twice daily for two doses on day one, then 200 mg twice daily on days two to five. Some patients with high cardiac risk were excluded. Select patients with severe COVID-19 and minimal cardiac risk also received azithromycin 500 mg x 1, then 250 mg daily on days two through five. Hydroxychloroquine was started within 48 hours of hospital admission in almost all patients. This study had several limitations. For example, the outcomes of almost 300 patients were not included in the analysis, and there were differences between treatment groups that could not be adequately adjusted for (e.g., baseline disease severity, other treatments received). • The FDA suggested, as part of their now-revoked Emergency Use Authorization for hydroxychloroquine, for hospitalized patients weighing ≥ 50 kg, a dose of 800 mg on day one, followed by 400 mg once daily for four to seven days.³¹ This suggested dosing regimen is unlikely to produce an antiviral effect.⁷³ • The WHO has discontinued the hydroxychloroquine arm of the Solidarity Trial because interim results suggest there is little mortality benefit for hospitalized patients.⁷⁴ • Outpatients (n=491) with confirmed or probable COVID-19 were randomized to hydroxychloroquine (800 mg x 1, then 600 mg six to eight hours later, then 600 mg once daily for four days) or placebo, starting within four days of symptom onset.¹¹ Mean age was 40 years, and 68% had no chronic medical conditions. Hydroxychloroquine was not better than placebo at improving symptom severity. At day 14, 24% of hydroxychloroquine patients were still symptomatic, vs 30% of placebo patients (p=0.21). Forty-three percent of hydroxychloroquine patients had side effects, vs 22% of placebo patients. Four hydroxychloroquine patients were hospitalized, and there was one outpatient death in this group. In the placebo group, ten placebo patients were hospitalized, one of which died (p=0.29). • Clinical trials are ongoing on the use of hydroxychloroquine to prevent COVID-19 in healthcare workers. • See www.clinicaltrials.gov for regimens being studied. <p>Safety</p> <ul style="list-style-type: none"> • In addition to efficacy concerns, the FDA's revocation of its EUA for hydroxychloroquine was based on adverse effects; its known and potential benefits no longer outweigh the known and potential side effects (e.g., serious cardiac events and other serious side effects).³³ • Due to the risk of arrhythmias, the FDA recommends against hydroxychloroquine use for COVID-19 outside of a <p><i>Continued...</i></p>

More...

Drug	Pertinent Information or Resources Note that DOSES provided are examples only for ADULTS ; the optimal dose has not been determined for any treatment.
Hydroxy-chloroquine, continued	<p>clinical trial.³³</p> <ul style="list-style-type: none"> • Adverse effects are not well-characterized at the doses studied for COVID-19. In general, potential adverse effects include: gastrointestinal side effects (take with food or milk), headache, hypoglycemia, QT prolongation and other conduction disturbances (especially with hypokalemia, hypomagnesemia, or heart disease), cardiomyopathy, myopathy, movement disorders, neurotoxicity, ocular toxicity, ototoxicity, anemia, thrombocytopenia, neutropenia, bone marrow suppression, serious dermatologic reactions, and psoriasis flare.^{27,31} Monitor electrolytes, glucose, complete blood count, electrocardiogram, baseline renal and hepatic function, knee and ankle reflexes, vision, and mental status.^{6,27,31} • When used with azithromycin (and other QT-prolonging medications), QT prolongation is of increased concern.^{2,6} Information on managing QT prolongation risk in these patients is available at https://www.ahajournals.org/doi/pdf/10.1161/CIRCULATIONAHA.120.047521.
IL-6 antagonist Tocilizumab (<i>Actemra</i>); sarilumab (<i>Kevzara</i>); siltuximab (<i>Sylvant</i>).	<ul style="list-style-type: none"> • Anti-IL-6 monoclonal antibody. • Some, but not all, data from China suggests an association between elevated IL-6 and severe COVID-19 disease.¹⁸ • Some anecdotal reports, case series, and cohort studies suggest benefit for tocilizumab (<i>Actemra</i>). One or two doses of 400 to 800 mg (4 to 8 mg/kg) has been used.^{18,68} • May cause increased infections, neutropenia, thrombocytopenia, and elevated liver enzymes.^{1,34-38} There are several cases of tocilizumab-associated worsening of COVID-19, perhaps due to immunosuppression, despite an associated reduction in inflammatory markers.⁸⁰ • Not for routine use. Clinical trials are planned or underway for treatment of pneumonia or cytokine storm. See www.clinicaltrials.gov for more information. • Outside of a clinical trial, limit to patients with evidence of cytokine storm (e.g., elevated IL-6, etc), with specialist consultation.⁴⁴ • The manufacturer of <i>Kevzara</i> (sarilumab) has discontinued its U.S. clinical trial in COVID-19 patients requiring mechanical ventilation (n = 194) because it did not meet its primary endpoint (improvement on a disease severity scale) or key secondary endpoints. The results of this study are not yet published.⁷⁷
Janus Kinase Inhibitors (Ruxolitinib [<i>Jakafi</i>], etc)	<ul style="list-style-type: none"> • No data. • Interest based on potential to block IL-6 effects, reduce cytotoxic T cells, and increase regulatory T cells. • NIH guidelines recommend against use, except in a clinical trial.⁵⁰ See www.clinicaltrials.gov for more information.

More . . .

Drug	Pertinent Information or Resources Note that DOSES provided are examples only for ADULTS ; the optimal dose has not been determined for any treatment.
Lopinavir/ ritonavir (Kaletra)	<ul style="list-style-type: none"> Lopinavir/ritonavir has not demonstrated anti-SARS-CoV-2 activity in humans.¹⁵ Small study suggested benefit (reduced composite endpoint of ARDS or death) for 2003 SARS vs historical control.¹⁷ Results from a randomized, open-label study (n=199) suggest it might reduce complications such as acute kidney injury, secondary infections, or need for mechanical ventilation in patients with COVID-19 pneumonia.¹⁵ However, time to clinical improvement was not reduced (main outcome measure).¹⁵ Gastrointestinal adverse effects may limit use.^{15,30} There is interest in studying lopinavir/ritonavir earlier in the disease course, or in combination with other medications.¹⁵ Use with ribavirin and interferon beta-1b early in the disease course (mean five days from symptom onset) was compared to lopinavir/ritonavir alone in hospitalized patients (n=127).⁵⁸ In this open-label study, median time to viral clearance was seven days with combination therapy vs 12 days for lopinavir/ritonavir alone.⁵⁸ Alleviation of symptoms occurred in four days vs eight days, respectively (p<0.0001).⁵⁸ The WHO has discontinued the lopinavir/ritonavir arm of the Solidarity Trial because interim results suggest no mortality benefit for hospitalized patients.⁷⁴ Additional clinical trials are planned or underway. See www.clinicaltrials.gov for more information.
Losartan, Telmisartan	<ul style="list-style-type: none"> Studies in mice suggest that ARBs can reduce lung damage caused by SARS-CoV.¹³ Clinical trials are underway for treatment of COVID-19. See www.clinicaltrials.gov for more information.
Oseltamivir	<ul style="list-style-type: none"> Not expected to be effective against SARS-CoV-2 because SARS-CoV-2 does not use neuraminidase.²⁶ Has been used for COVID-19 pneumonia, but there is no efficacy data.¹²
Remdesivir	<ul style="list-style-type: none"> Remdesivir has <i>in vitro</i> activity against SARS-CoV-2.⁴⁰ In a cohort of 53 evaluable patients treated with remdesivir for severe COVID-19 disease, use was associated with clinical improvement in regard to oxygen support requirements in 68% of patients.⁴⁰ Mortality was 13%, which is less than in other case series and cohorts.⁴⁰ The most common adverse events were liver enzyme elevation (23%), diarrhea (9%), rash, renal impairment, hypotension (8%), acute kidney injury, atrial fibrillation, multiorgan dysfunction, hypernatremia, and venous thrombosis (6%).⁴⁰ Causality could not be assessed due to the effects of COVID-19 itself.⁴⁰ Based on previous data, mild to moderate transaminase elevations are expected with remdesivir.⁴⁰ Viral load was not evaluated,⁴⁰ but in a previous case report, virologic improvement was seen.⁸ Preliminary analysis of a double-blind, placebo-controlled trial (n = 1,063), remdesivir seemed to shorten time to recovery (11 days vs 15 days; p <0.001), but mortality was not statistically different (8% vs 11.6%; p = 0.059).^{54,72} Similarly, a Chinese study found a nonsignificant trend toward faster recovery.⁶¹ Unpublished data from the open-label SIMPLE-Severe study compared remdesivir-treated patients (n=312) to a matched cohort of patients receiving standard care (n=818). About 74% of remdesivir patients recovered by day 14 vs 59% of the
<i>Continued...</i>	

More...

Drug	Pertinent Information or Resources
Remdesivir, continued	<p>Note that DOSES provided are examples only for ADULTS; the optimal dose has not been determined for any treatment.</p> <p>standard-care patients. Mortality rate at day 14 was 7.6% in the remdesivir patients vs 12.5% in the standard-treatment group (OR 0.38, 95% CI 0.22 to 0.68, p = 0.001).⁷⁹</p> <ul style="list-style-type: none"> • U.S.: Remdesivir is being distributed to select hospitals by the government through Emergency Use Authorization.⁵⁷ This is a rapidly changing situation. For other potential opportunities for availability, see www.clinicaltrials.gov, www.gilead.com/remdesivir, or contact Gilead at 833-445-3230 (GILEAD-0) or GileadClinicalTrials@gilead.com. <ul style="list-style-type: none"> • The FDA has a fact sheet on remdesivir, including criteria for use, adverse effects, dosing, and more (https://www.fda.gov/media/137566/download). • Canada: Remdesivir is available through an Expanded Access Treatment Protocol at approved clinical trial sites. A list of active sites is available at https://www.canada.ca/en/health-canada/services/drugs-health-products/covid19-clinical-trials/list-authorized-trials.html and at www.clinicaltrials.gov. At this point, Gilead is not accepting more sites for consideration. Compassionate use requests continue to be reviewed for pregnant women and children <18 years of age with confirmed severe COVID-19.⁵⁹ • If remdesivir availability is limited at your hospital, the NIH recommends prioritizing use for patients requiring oxygen, but not high-flow oxygen, noninvasive ventilation, mechanical ventilation, or ECMO.⁵⁰ • Coadministration of remdesivir and chloroquine or hydroxychloroquine is not recommended based on <i>in vitro</i> data showing that these drugs might interfere with the metabolic activation and antiviral activity of remdesivir.⁵³ In Simple-Severe, recovery rate at day 14 for patients who received hydroxychloroquine plus remdesivir was lower than in patients who received remdesivir alone (57% percent vs 69%, HR 0.61, 95% CI 0.45 to 0.83, p=0.002). Concomitant hydroxychloroquine use was not associated with increased mortality, but was associated with a higher risk of adverse events.⁷⁹ • Another potential drug interaction involves inhibition of remdesivir elimination from hepatocytes by P-glycoprotein inhibitors. This interaction could result in hepatotoxicity.⁷⁶
Ribavirin	<ul style="list-style-type: none"> • Not potent enough to be effective at safe doses; hematologic toxicity precludes use.²⁶ See lopinavir/ritonavir section for information on combination use.
Statins	<ul style="list-style-type: none"> • No data. • Interest based on cardiovascular damage noted in COVID-19 patients and anti-inflammatory effects. Simvastatin might also block viral cell entry. • NIH guidelines recommend against use specifically for COVID-19 treatment outside of a clinical trial.⁵⁰ • See www.clinicaltrials.gov for more information on planned or ongoing studies.
tPA (alteplase)	<ul style="list-style-type: none"> • No data. • Interest based on reports of hypercoagulability in COVID-19 patients.¹⁹ • Studies underway to treat ARDS in COVID-19 patients.¹⁹

More . . .

Drug	Pertinent Information or Resources Note that DOSES provided are examples only for ADULTS ; the optimal dose has not been determined for any treatment.
Vaccines	<ul style="list-style-type: none"> • Due to evidence of a non-specific protective effect against respiratory infections, BCG vaccine is being studied to prevent COVID-19 disease in healthcare workers. See www.clinicaltrials.gov for more information. • Oral polio vaccine has been mentioned on the internet, but no trials are planned at this time.
Vitamin C	<ul style="list-style-type: none"> • Intravenous vitamin C is being studied for treatment of severe COVID-19 disease based on previous data in sepsis and ARDS. However, there is no clear evidence of benefit even for these conditions.⁴⁸ • Oral vitamin C is being studied for treatment of COVID-19 disease in the outpatient setting, and as prophylaxis. • See www.clinicaltrials.gov for more information on these planned or ongoing studies.
Vitamin D	<ul style="list-style-type: none"> • There is false information circulating that vitamin D is recommended by health officials. Interest in vitamin D stems from its effects on the immune system and pulmonary ACE2 expression. Studies are planned or underway using vitamin D for prevention or as a treatment adjunct. See www.clinicaltrials.gov for more information.
Zinc	<ul style="list-style-type: none"> • Zinc has <i>in vitro</i> activity against SARS-CoV.⁴⁷ • Studies of oral zinc, alone or in combination (e.g., with vitamin C, vitamin D, hydroxychloroquine [purported to help zinc get inside the cells⁴⁷], azithromycin) to prevent COVID-19 disease are planned or ongoing. • See www.clinicaltrials.gov for more information.

—Continue to the next page for information to help answer patient COVID-19 questions and address misinformation—

More. . .

FAQs ABOUT COVID-19 AND PHARMACOTHERAPY

There is a lot of misinformation regarding COVID-19 on the internet. Use this table to help answer patient questions and correct misconceptions.

Clinical question	Pertinent information or resource
Do ACE inhibitors or ARBs make COVID-19 worse?	<ul style="list-style-type: none"> • The SARS-CoV-2 virus uses ACE2 to enter cells.¹³ ACE inhibitors and ARBs may upregulate ACE2.¹³ In theory, these drugs could thereby facilitate virus entry into cells.¹³ But on the other hand, blocking angiotensin could reduce lung injury.¹³ • No evidence suggests that patients taking an ACEI or ARB are more susceptible to COVID-19 infection, or that these medications worsen outcomes.^{14,51,52,69} One cohort study even suggests reduced mortality in COVID-19 patients taking them for other indications.⁵¹ Furthermore, we know that these drugs benefit patients with diabetic nephropathy and cardiovascular disease, populations at risk of severe COVID-19 disease.^{13,22} • Patients should continue these medications. See statements from: <ul style="list-style-type: none"> • the American Heart Association, the Heart Failure Society of America, and the American College of Cardiology at https://newsroom.heart.org/news/patients-taking-ace-i-and-arbs-who-contract-covid-19-should-continue-treatment-unless-otherwise-advised-by-their-physician. • the Canadian Cardiovascular Society at https://www.ccs.ca/images/Images_2020/CCS_CHFS_Update_COVID__CV_medications_Mar20.pdf. • the European Society of Cardiology at https://www.escardio.org/Councils/Council-on-Hypertension-(CHT)/News/position-statement-of-the-esc-council-on-hypertension-on-ace-inhibitors-and-ang.
Can NSAIDs be used in COVID-19-infected patients?	<ul style="list-style-type: none"> • Anecdotal reports regarding worse COVID-19 outcomes in patients taking NSAIDs have spread in the media and on social media, including via a tweet from a French health official.^{14,23} In 2019, a French report suggested that NSAIDs could worsen infections, mainly Strep, perhaps by masking symptoms.^{24,25} However, there is currently no reliable clinical data supporting worse outcomes in patients taking NSAIDs or aspirin.^{14,20} Preclinical data is mixed on the potential effects of NSAIDs on COVID-19 (increased expression of ACE2, which the virus uses to enter cells, vs potential antiviral activity of NSAIDs).¹⁴ • Patients taking low-dose aspirin should not stop taking it because of COVID-19 concerns.¹⁴ • Neither the FDA nor Health Canada is advising changes to NSAID use due to COVID-19.^{20,21}
Are any supplements effective for prevention or treatment of COVID-19?	<ul style="list-style-type: none"> • There is no scientific evidence that any alternative remedies can prevent or treat COVID-19, and some products may not be safe.⁵ See our <i>Natural Medicines</i> database (www.naturaldatabase.com) for information on efficacy and safety of specific alternative medicines. • A study using honey as an adjunct to standard care for treatment of COVID-19 is planned. • Several studies are looking at multivitamin/mineral combos as adjuncts for treatment or prevention. See www.clinicaltrials.gov for more information. For information on zinc, vitamin C, and vitamin D see the “Treatments of Interest” chart, above.

More . . .

Clinical question	Pertinent information or resource
Are heartburn drugs effective for treating or preventing COVID-19?	<ul style="list-style-type: none"> • For information on famotidine, see the “Treatments of Interest” chart, above. • Proton pump inhibitor (PPI) use was associated with a higher risk of COVID-19 based on results of an online survey of ~53,000 English-speaking Americans, after controlling for confounders. Risk was higher for patients taking a PPI twice daily vs once daily. H2 blockers did not appear to increase risk. Any increased absolute risk of COVID-19 in PPI users is probably small. If patients ask about PPI use, consider this an opportunity to reevaluate treatment and reinforce COVID-19 preventive measures.⁷⁸
Does nicotine protect against COVID-19?	<ul style="list-style-type: none"> • In China, there was an unexpectedly low prevalence of smoking among patients hospitalized with COVID-19. Low smoking prevalence among hospitalized COVID-19 patients has also been seen in the U.S.⁵⁶ • Nicotine, through its cholinergic agonist activity, blocks production of inflammatory cytokines such as IL-6.⁵⁶ • There is interest in using nicotine, either as currently available products, or perhaps via nebulization, as an adjunct for COVID-19 treatment.⁵⁶ • Continue to use nicotine replacement products for nicotine users who are hospitalized for COVID-19, and for anyone who desires to quit smoking.⁵⁶

Users of this resource are cautioned to use their own professional judgment and consult any other necessary or appropriate sources prior to making clinical judgments based on the content of this document. Our editors have researched the information with input from experts, government agencies, and national organizations. Information and internet links in this article were current as of the date of publication.

More . . .

Levels of Evidence

In accordance with our goal of providing Evidence-Based information, we are citing the **LEVEL OF EVIDENCE** for the clinical recommendations we publish.

Level	Definition	Study Quality
A	Good-quality patient-oriented evidence.*	<ol style="list-style-type: none"> 1. High-quality RCT 2. SR/Meta-analysis of RCTs with consistent findings 3. All-or-none study
B	Inconsistent or limited-quality patient-oriented evidence.*	<ol style="list-style-type: none"> 1. Lower-quality RCT 2. SR/Meta-analysis with low-quality clinical trials or of studies with inconsistent findings 3. Cohort study 4. Case control study
C	Consensus; usual practice; expert opinion; disease-oriented evidence (e.g., physiologic or surrogate endpoints); case series for studies of diagnosis, treatment, prevention, or screening.	

***Outcomes that matter to patients** (e.g., morbidity, mortality, symptom improvement, quality of life).

RCT = randomized controlled trial; **SR** = systematic review

[Adapted from Ebell MH, Siwek J, Weiss BD, et al. Strength of Recommendation Taxonomy (SORT): a patient-centered approach to grading evidence in the medical literature. *Am Fam Physician* 2004;69:548-56. <http://www.aafp.org/afp/2004/0201/p548.pdf>.]

Project Leader in preparation of this clinical resource (360423): *Melanie Cupp, Pharm.D., BCPS; Last modified July 27, 2020.*

References

1. Product monograph for *Kevzara*. Sanofi Genzyme. Mississauga, ON L4W 4V9. August 2019.
2. Gautret P, Lagier JC, Parola P, et al. Hydroxychloroquine and azithromycin as a treatment of COVID-19: results of an open-label non-randomized clinical trial. *Int J Antimicrob Agents* 2020 Mar 20:105949. doi: 10.1016/j.ijantimicag.2020.105949.
3. Gao J, Tian Z, Yang X. Breakthrough: chloroquine phosphate has shown apparent efficacy in treatment of COVID-19 associated pneumonia in clinical studies. *BioSci Trends* 2020;14:72-3.
4. FDA. Fact sheet for health care providers emergency use authorization (EUA) of chloroquine phosphate supplied from the strategic national stockpile for treatment of COVID-19 in certain hospitalized patients. Revoked. June 15, 2020. <https://www.fda.gov/media/136535/download>. (Accessed April 15, 2020).
5. NIH. National Center for Complementary and Integrative Health. In the news: coronavirus and "alternative" treatments. <https://www.nccih.nih.gov/health/in-the-news-coronavirus-and-alternative-treatments>. (Accessed April 15, 2020).
6. Cortegiani A, Ingoglia G, Ippolito M, et al. A systematic review on the efficacy and safety of chloroquine for the treatment of COVID-19. *J Crit Care* 2020 Mar 10. doi: 10.1016/j.jcrr.2020.03.005.
7. Bermejo-Martin JF, Kelvin DJ, Eiros JM, et al. Macrolides for the treatment of severe respiratory illness caused by novel H1N1 swine influenza viral strains. *J Infect Dev Ctries* 2009;3:159-61.
8. Ko WC, Rolain JM, Lee NY, et al. Arguments in favour of remdesivir for treating SARS-CoV-2 infections. *Int J Antimicrob Agents* 2020 Mar 6:105933. doi: 10.1016/j.ijantimicag.2020.105933.
9. Deftereos SG, Giannopoulos G, Vrachatis DA, et al. Effect of colchicine vs standard care on cardiac and inflammatory biomarkers and clinical outcomes in patients hospitalized with coronavirus disease 2019: the GRECCO-19 randomized clinical trial. *JAMA Netw Open* 2020;3:e2013136.
10. World Health Organization. Clinical management of severe acute respiratory infection when COVID-19 is suspected. Interim guidance. May 27, 2020. [https://www.who.int/publications-detail/clinical-management-of-severe-acute-respiratory-infection-when-novel-coronavirus-\(ncov\)-infection-is-suspected](https://www.who.int/publications-detail/clinical-management-of-severe-acute-respiratory-infection-when-novel-coronavirus-(ncov)-infection-is-suspected). (Accessed July 18, 2020).
11. Skipper CP, Pastick KA, Engen NW, et al. Hydroxychloroquine in nonhospitalized adults with early COVID-19. *Ann Intern Med* 2020 Jul 16. doi.org/10.7326/M20-4207.
12. Chen N, Zhou M, Dong X, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet* 2020;395:507-13.
13. Kuster GM, Pfister O, Burkard T, et al. SARS-CoV2: should inhibitors of the renin-angiotensin system be withdrawn in patients with COVID-19? *Eur Heart J* 2020 Mar 20. doi: 10.1093/eurheartj/ehaa325.
14. Canadian Cardiovascular Society. Guidance from the CCS COVID-19 Rapid Response team. March 20, 2020. https://www.ccs.ca/images/Images_2020/CCS_CHF_S_Update_COVID_CV_medications_Mar20.pdf. (Accessed April 15, 2020).
15. Cao B, Wang Y, Wen W, et al. A trial of lopinavir-ritonavir in adults hospitalized with severe COVID-19. *N Engl J Med* 2020 Mar 18. doi: 10.1056/NEJMoa2001282.

More . . .

16. Wu C, Chen X, Cai Y, et al. Risk factors associated with acute respiratory distress syndrome and death in patients with coronavirus disease 2019 pneumonia in Wuhan, China. *JAMA Intern Med* 2020 Mar 13. doi: 10.1001/jamainternmed.2020.0994.
17. Chu CM, Cheng VC, Hung IF, et al. Role of lopinavir/ritonavir in the treatment of SARS: initial virological and clinical findings. *Thorax* 2004;59:252-6.
18. Matos RI, Chung KK. DoD COVID-19 practice management guide. Clinical management of COVID-19. March 23, 2020.
19. Anon. New treatment approach proposed for respiratory distress in COVID-19. *Drug Topics*. March 26, 2020. <https://www.drugtopics.com/clinical-news/new-treatment-approach-proposed-respiratory-distress-covid-19>. (Accessed April 15, 2020).
20. FDA. FDA advises patients on use of non-steroidal anti-inflammatory drugs (NSAIDs) for COVID-19. March 19, 2020. <https://www.fda.gov/drugs/drug-safety-and-availability/fda-advises-patients-use-non-steroidal-anti-inflammatory-drugs-nsaids-covid-19>. (Accessed April 15, 2020).
21. Health Canada. No scientific evidence that ibuprofen worsens COVID-19 symptoms. March 20, 2020. <https://healthycanadians.gc.ca/recall-alert-rappel-avis/hc-sc/2020/72633a-eng.php>. (Access April 15, 2020).
22. Yang X, Yu Y, Xu J, et al. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respir Med* 2020 Feb 24. doi: 10.1016/S2213-2600(20)30079-5.
23. Melville NA, Nainggolan L. Are warnings against NSAIDs in COVID-19 warranted? March 17, 2020. <https://www.medscape.com/viewarticle/926940>. (Accessed April 15, 2020).
24. European Medicines Agency. EMA gives advice on the use of non-steroidal anti-inflammatories for COVID 19. March 18, 2020. <https://www.ema.europa.eu/en/news/ema-gives-advice-use-non-steroidal-anti-inflammatories-covid-19>. (Accessed April 15, 2020).
25. Agence Nationale de Securite du Medicament et des Produits de Sante. Anti-inflammatoires non steroïdiens (AINS) et complications infectieuses graves-point d'information. April 18, 2019. <https://www.ansm.sante.fr/S-informer/Points-d-information-Points-d-information/Anti-inflammatoires-non-steroidiens-AINS-et-complications-infectieuses-graves-Point-d-Information>. (Accessed April 15, 2020).
26. McCreary EK, Pogue JM. COVID-19 treatment: a review of early and emerging options. *Open Forum Infectious Diseases*. *Open Forum Infect Dis* 2020 March 23;7(3): ofaa105. doi: 10.1093/ofid/ofaa105.
27. Clinical Pharmacology powered by ClinicalKey. Tampa (FL): Elsevier. 2020. <http://www.clinicalkey.com>. (Accessed April 15, 2020).
28. Gautret P, Lagier JC, Parola P et al. Clinical and microbiological effect of a combination of hydroxychloroquine and azithromycin in 80 COVID-19 patients with at least a six-day follow-up: an observational study. <https://www.mediterranee-infection.com/wp-content/uploads/2020/03/COVID-IHU-2-1.pdf>. (Accessed April 15, 2020).
29. Chen J, Danping L, Li L, et al. A pilot study of hydroxychloroquine in treatment of patients with common coronavirus disease-19 (COVID-19). *J Zhejiang Univ* 2020;49. doi: 10.3785/j.issn. 1008-9292.2020.03.03.
30. Young BE, Ong SWX, Kalimuddin S, et al. Epidemiologic features and clinical course of patients infected with SARS-CoV-2 in Singapore. *JAMA* 2020 Mar 3. doi: 10.1001/jama.2020.3204.
31. RECOVERY Collaborative Group. Dexamethasone in hospitalized patients with COVID-19-preliminary report. *N Engl J Med* 2020 Jul 17. doi:10.1056/NEJMoa2021436.
32. CDC. Information for clinicians on therapeutic options for COVID-19 patients. April 13, 2020. <https://www.cdc.gov/coronavirus/2019-ncov/hcp/therapeutic-options.html>. (Accessed April 15, 2020).
33. FDA. Frequently asked questions on the revocation of the Emergency Use Authorization for hydroxychloroquine sulfate and chloroquine phosphate. June 19, 2020. <https://www.fda.gov/media/138946/download>. (Accessed July 1, 2020).
34. Product information for *Kevzara*. Sanofi-Aventis. Bridgewater, NJ 08807. April 2018.
35. Product information for *Actemra*. Genentech. South San Francisco, CA 94080. June 2019.
36. Product information for *Sylvant*. EUSA Pharma. Hertfordshire, U.K. HP2 4TZ. December 2019.
37. Product monograph for *Actemra*. Mississauga, ON L5N 5M8. September 2019.
38. Product monograph for *Sylvant*. Janssen. Toronto, ON M3C 1L9. March 2018.
39. Chen Z, Hu JM, Zhang Z, et al. Efficacy of hydroxychloroquine in patients with COVID-19: results of a randomized clinical trial. medRxiv 2020.03.22.20040758. <https://doi.org/10.1101/2020.03.22.20040758>. (Accessed April 15, 2020).
40. Grein J, Ohmagari N, Shin D, et al. Compassionate use of remdesivir for patients with severe COVID-19. *N Engl J Med* 2020 Apr 10. doi: 10.1056/NEJMoa2007016.
41. Borba MG, Val FF, Sampaio VS, et al. Effect of high vs low doses of chloroquine diphosphate as adjunctive therapy for patients hospitalized with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection: a randomized clinical trial. *JAMA Netw Open* 2020;3(4.23):e208857. doi:10.1001/jamanetworkopen.2020.8857.
42. Tang W, Cao Z, Han M, et al. Hydroxychloroquine in patients with mainly mild to moderate coronavirus disease 2019: open-label, randomized, controlled trial. *BMJ* 2020;369. doi: 10.1136/bmj.m1849.

More . . .

43. Mahevas M, Tran VT, Roumier M, et al. No evidence of clinical efficacy of hydroxychloroquine in patients hospitalized for COVID-19 infection with oxygen requirement: results of a study using routinely collected data to emulate a target trial. medRxiv 2020.04.10.20060699. <https://doi.org/10.1101/2020.04.10.20060699>. (Accessed April 16, 2020).
44. British Columbia Ministry of Health. Unproven therapies for COVID-19. Updated July 14, 2020. http://www.bccdc.ca/Health-Professionals-Site/Documents/Guidelines_Unproven_Therapies_COVID-19.pdf. (Accessed July 27, 2020).
45. Mehta P, McAuley DF, Brown M, et al. COVID-19: consider cytokine storm syndromes and immunosuppression. *Lancet* 2020;395:1033-4.
46. Infectious Diseases Society of America guidelines on the treatment and management of patients with COVID-19. Last updated June 25, 2020. <https://www.idsociety.org/practice-guideline/covid-19-guideline-treatment-and-management/>. (Accessed July 1, 2020).
47. te Velthuis AJ, van den Worn SH, Sims AC, et al. Zn(2+) inhibits coronavirus and arterivirus RNA polymerase activity in vitro and zinc ionophores block the replication of these viruses in cell culture. *PLoS Pathog* 2010;6(11):e1001176. doi: 10.1371/journal.ppat.1001176.
48. Surani S, Sharma M. Revisiting the role of vitamin C in sepsis. Is it a forlorn hope or is there still dearth of data? *Open Respir Med J* 2019;13:55-7.
49. Magagnoli J, Narendran S, Pereira F, et al. Outcomes of hydroxychloroquine usage in United States veterans hospitalized with Covid-19. medRxiv. 2020.04.16.20065920. <https://doi.org/10.1101/2020.04.16.20065920>. (Accessed April 16, 2020).
50. NIH. Coronavirus disease 2019 (COVID-19). Treatment guidelines. Last updated July 24, 2020. <https://covid19treatmentguidelines.nih.gov/>. (Accessed July 27, 2020).
51. Zhang P, Zhu L, Cai J, et al. Association of inpatient use of angiotensin converting enzyme inhibitors and angiotensin II receptor blockers with mortality among patients with hypertension hospitalized with COVID-19. *Circ Res* 2020 Apr 17. doi: 10.1161/CIRCRESAHA.120.317134.
52. Li J, Wang X, Chen J, et al. Association of renin-angiotensin system inhibitors with severity or risk of death in patients with hypertension hospitalized for coronavirus disease 2019 (COVID-19) infection in Wuhan, China. *JAMA Cardiol* 2020 Apr 23. doi:10.1001/jamacardio.2020.1624.
53. FDA. Fact sheet for health care providers emergency use authorization (EUA) of remdesivir (GS-5734). June 2020. <https://www.fda.gov/media/137566/download>. (Accessed June 16, 2020).
54. FDA. Fact sheet for health care providers. Emergency use authorization (EUA) for remdesivir (GS-5734). <https://www.fda.gov/media/137566/download>. (Accessed May 1, 2020).
55. Rogosnitzky M, Berkowitz E, Jadad AR. Delivering benefits at speed through real-world repurposing of off-patent drugs: the COVID-19 pandemic as a case point. *JMIR Public Health Surveill* 2020;6:e19199.
56. Farsalinos K, Barbouni A, Niaura R. Systematic review of the prevalence of current smoking among hospitalized COVID-19 patients in China: could nicotine be a therapeutic option? *Intern Emerg Med* 2020 May 9. doi: 10.1007/s11739-020-02355-7.
57. Ison MG, Wolfe C, Boucher HW. Emergency use authorization of remdesivir: the need for a transparent distribution process. *JAMA* 2020 May 14. doi: 10.1001/jama.2020.8863.
58. Hung IF, Lung KC, Tso EY, et al. Triple combination of interferon beta-1b, lopinavir-ritonavir, and ribavirin in the treatment of patients admitted to hospital with COVID-19: an open-label, randomized, phase 2 trial. *Lancet* 2020 May 8. doi: 10.1016/S0140-6736(20)31042-4.
59. Personal communication (written). Lisa Bulet. Medical Information. Gilead Sciences Canada, Inc. Mississauga, ON L5N 2W3. May 14, 2020.
60. Geleris J, Sun Y, Platt J, et al. Observational study of hydroxychloroquine in hospitalized patients with COVID-19. *N Engl J Med* 2020 May 7. doi: 10.1056/NEJMoa2012410.
61. Wang Y, Zhang D, Du G, et al. Remdesivir in adults with severe COVID-19: a randomised, double-blind, placebo-controlled, multicentre trial. *Lancet* 2020 April 29. doi: 10.1016/S0140-6736(20)31022-9.
62. Shen C, Wang Z, Zhao F, et al. Treatment of 5 critically ill patients with COVID-19 with convalescent plasma. *JAMA* 2020;323:1582-9.
63. Duan K, Liu B, Li C, et al. Effectiveness of convalescent plasma therapy in severe COVID-19 patients. *Proc Natl Acad Sci USA* 2020;117:9490-6.
64. Ye M, Fu D, Ren Y, et al. Treatment with convalescent plasma for COVID-19 patients in Wuhan, China. *J Med Virol* 2020 Apr 15. doi: 10.1002/jmv.25882.
65. Cavalli G, De Luca G, Campochiaro C, et al. Interleukin-1 blockade with high-dose anakinra in patients with COVID-19, acute respiratory distress syndrome, and hyperinflammation: a retrospective cohort study. *Lancet Rheumatol* 2020 May 7. doi: 10.1016/S2665-9913(20)30127-2.
66. Rosenberg ES, Dufort EM, Udo T, et al. Association of treatment with hydroxychloroquine or azithromycin with in-hospital mortality in patients with COVID-19 in New York State. *JAMA* 2020 May 11. doi: 10.1001/jama.2020.8630.
67. Freedberg DE, Conigliaro J, Wang TC, et al. Famotidine use is associated with improved clinical outcomes in hospitalized COVID-19 patients: a propensity score matched retrospective cohort study. *Gastroenterology* 2020 May 22. doi: 10.1053/j.gastro.2020.05.053.

More . . .

68. Xu X, Han M, Li T, et al. Effective treatment of severe COVID-19 patients with tocilizumab. *Proc Natl Acad Sci USA* 2020 Apr 29. doi: 10.1073/pnas.2005615117.
69. Reynolds HR, Adhikari S, Pulgarin C, et al. Renin-angiotensin-aldosterone system inhibitors and risk of COVID-19. *N Engl J Med* 2020 May 1. doi: 10.1056/NEJMoa2008975.
70. Rubin R. Testing an old therapy against a new disease: convalescent plasma for COVID-19. *JAMA* 2020 Apr 30. doi: 10.1001/jama.2020.7456.
71. Expanded Access protocol. Convalescent plasma for the treatment of patients with COVID-19. NCT 04338360. May 4, 2020. <https://www.uscovidplasma.org/pdf/COVID-19%20Plasma%20EAP.pdf>. (Accessed May 16, 2020).
72. Beigel JH, Tomashek KM, Dodd LE, et al. Remdesivir for the treatment of COVID-19—preliminary report. *N Engl J Med* 2020 May 22. doi: 10.1056/NEJMoa2007764.
73. FDA. Letter to Gary L. Disbrow, Deputy Assistant Secretary, Director, Medical Countermeasures Programs, BARDA, in response to request for revocation of Emergency Use Authorization for chloroquine phosphate and hydroxychloroquine. June 15, 2020. <https://www.fda.gov/media/138945/download>. (Accessed June 15, 2020).
74. Bagozzi D. WHO discontinued hydroxychloroquine and lopinavir/ritonavir treatment arms for COVID-19. News release. July 4, 2020. <https://www.who.int/news-room/detail/04-07-2020-who-discontinues-hydroxychloroquine-and-lopinavir-ritonavir-treatment-arms-for-covid-19>. (Accessed July 6, 2020).
75. Arshad S, Kilgore P, Chaudry ZS, et al. Treatment with hydroxychloroquine, azithromycin, and combination in patients hospitalized with COVID-19. *Int J Infect Dis* 2020 Jul 1:S1201-9712(20)30534-8. doi: 10.1016/j.ijid.2020.06.099.
76. Leegwater E, Strik A, Wilms EB, et al. Drug induced liver injury in a COVID-19 patient: potential interaction of remdesivir with P-glycoprotein inhibitors. *Clin Infect Dis* 2020 Jun 28:ciaa883. doi: 10.1093/cid/ciaa883.
77. Anon. Sanofi and Regeneron provide update on Kevzara (sarilumab) phase 3 U.S. trial in COVID-19 patients. July 2, 2020. <http://www.news.sanofi.us/2020-07-02-Sanofi-and-Regeneron-provide-update-on-Kevzara-R-sarilumab-Phase-3-U-S-trial-in-COVID-19-patients>. (Accessed July 8, 2020).
78. American College of Gastroenterology. Information sheet and FAQs about proton pump inhibitors (PPIs) and risk of COVID-19. July 7, 2020. https://webfiles.gi.org/links/media/ACG_Almario_et_al_Info_Sheet_and_FAQs_About_PPIs_COVID19_07_072020_FINAL.pdf. (Accessed July 9, 2020).
79. Gilead press release. Gilead presents additional data on investigational antiviral remdesivir for the treatment of COVID-19. July 10, 2020. <https://www.gilead.com/news-and-press/press-room/press-releases/2020/7/gilead-presents-additional-data-on-investigational-antiviral-remdesivir-for-the-treatment-of-covid-19>. (Accessed July 11, 2020).
80. Radbel J, Narayanan N, Bhatt PJ. Use of tocilizumab for COVID-19-induced cytokine release syndrome: a cautionary case report. *Chest* 2020;158:e15-9.

Cite this document as follows: Clinical Resource, COVID-19 and Pharmacotherapy. Pharmacist's Letter/Prescriber's Letter. April 2020.

trc | pharmacist's letter™

Evidence and Recommendations You Can Trust...

trc | prescriber's letter™

trc | pharmacy technician's letter™

3120 West March Lane, Stockton, CA 95219 ~ TEL (209) 472-2240 ~ FAX (209) 472-2249

Copyright © 2020 by Therapeutic Research Center

Subscribers to the *Letter* can get clinical resources, like this one, on any topic covered in any issue by going to pharmacist.therapeuticresearch.com ~ prescriber.therapeuticresearch.com ~ pharmacytech.therapeuticresearch.com